




HOT DAYS ALONG THE WEST ANTARCTIC PENINSULA



LTER Palmer has maintained a 19 year time series along the West Antarctic Peninsula

The Boss! **Current team**



PI Hugh Ducklow (MBL)
Bacteria-Biogeochemistry



Bill Fraser (Polar Associates)
- Penguins & Fish



Karen Baker (Scripps)
- Data management
& Informatics



Scott Doney (WHOI)
- Ocean Modeling



Beth Simmons (Scripps)
- Education &
Outreach




Oscar Schofield (Rutgers) - Phytoplankton
Doug Martinson (LDEO) - Ocean Physics
Debbie Steinberg (VIMS) - Zooplankton



Sharon Stammerjohn (UCSC)
- Climate and Ice

Our Current grid



Acknowledgements to past LTER PIs: Ray Smith, Barbara Prezelin, Robin Ross, Langdon Quetin, Dave Karl, Maria Vernet, Eileen Hoffman, John Klinck, Dave Karl



The man!

Increases in Antarctic penguin populations: reduced competition with whales or a loss of sea ice due to environmental warming?

William R. Fraser^{*,**}, Wayne Z. Trivelpiece^{*}, David G. Ainley and Susan G. Trivelpiece^{*}

Point Reyes Bird Observatory, 4990 Shoreline Highway, Stinson Beach, CA 94970, USA

Received 25 June 1991; accepted 6 August 1991

Summary. A central tenet of Antarctic ecology suggests that increases in Chinstrap Penguin (*Pygoscelis antarctica*) populations during the last four decades resulted from an increase in prey availability brought on by the decrease in baleen whale stocks. We question this tenet and present evidence to support the hypothesis that these increases are due to a gradual decrease in the frequency of cold years with extensive winter sea ice cover resulting from environmental warming. Supporting data were derived from one of the first, major multidisciplinary winter expedition to the Scotia and Weddell seas; recent satellite images of ocean ice cover; and the analysis of long-term surface temperature records and penguin demography. Our observations indicate there is a need to pay close attention to environmental data in the management of Southern Ocean resources given the complexity of relating biological changes to ecological perturbations.

Introduction

Populations of many krill-eating, Southern Ocean predators have exhibited significant changes during the last four decades. Notable among these, have been increases in the abundance of Chinstrap Penguins (*Pygoscelis antarctica*), which breed mainly on the Antarctic Peninsula and islands of the Scotia Sea (Watson 1975). At many colonies, numbers have increased 6–10% per annum (Laws 1985), and at some localities fivefold increases have occurred in the last 20 years (Rootes 1988). Chinstraps have also expanded their range southward along the western side of the Antarctic Peninsula (Parmelee and Parmelee 1987; Poncet and Poncet 1987) into areas historically dominated by the closely related adelic Penguin (*P. adeliae*; Fig. 1). A central tenet of Antarctic ecology ex-

plains these population changes in terms of a presumed increase in food availability that resulted from the decrease in baleen whale stocks due to commercial whaling (Sladen 1964; Emison 1968; Conroy 1975; Croxall and Kirkwood 1979; Croxall and Prince 1979; Croxall et al. 1984). This tenet is based on the fact that the dominant component in the summer diets of both Chinstraps and whales is the Antarctic krill (*Euphausia superba*). Although this tenet has been widely accepted, the possible mechanism by which a decrease in whales could have led to an increase in Chinstraps has not been questioned (cf. Horwood 1980). Indeed, the long-standing view has simply been that whaling led to a "krill surplus" that was used by krill-eating predators when competitive release altered the existing patterns of consumption (Laws 1985).

Although this whale reduction hypothesis has clearly been useful in guiding research on trophic interactions in the Southern Oceans, it is now apparent that increases in Chinstrap populations have not been mirrored by their sympatric, most closely related congener, the Adelic Penguin. Adelines share a significant portion of their range on the Antarctic peninsula and islands of the Scotia Arc with Chinstraps (Watson 1975). Alike in size and general appearance, both exhibit broad ecological similarities, not the least of which is a predominance of krill in their summer diets (Volkman et al. 1980; Trivelpiece et al. 1987, 1990; Trivelpiece and Trivelpiece 1990). Yet, when compared to Chinstraps, population increases in Adelines have not been as substantial, and at many sites appear to represent nothing more than recovery after human disturbance and exploitation (Poncet and Poncet 1987). Adelines, in fact, have declined noticeably at several localities on the Antarctic Peninsula, a change considered "unexplainable" by Poncet and Poncet (1987). This raises an interesting challenge to the whale reduction hypothesis: If the decrease of baleen whale stocks actually led to a krill surplus, why have populations of the ecologically similar Adelines residing in the same geographical areas shown such different responses?

Here we propose that the answer to this question does not rest with the idea of a krill surplus. Instead, we suggest

Key point: If the decimation of baleen whale populations did in fact lead to a "krill surplus", why were krill-dependent, top predator populations exhibiting such dichotomous trends?

* Current address: Polar Oceans Research Group, Department of Oceanography, Old Dominion University, Norfolk, VA 23529, USA.

** Present address: W.R. Fraser, ODU Central States Office 830 Hunt Farm Rd., Long Lake, MN 55356, USA

Offprint requests to: W.R. Fraser



The man!

Increases in Antarctic penguin populations: reduced competition with whales or a loss of sea ice due to environmental warming?

William R. Fraser^{*,**}, Wayne Z. Trivelpiece^{*}, David G. Ainley and Susan G. Trivelpiece^{*}

Point Reyes Bird Observatory, 4990 Shoreline Highway, Stinson Beach, CA 94970, USA

Received 25 June 1991; accepted 6 August 1991

Summary. A central tenet of Antarctic ecology suggests that increases in Chinstrap Penguin (*Pygoscelis antarctica*) populations during the last four decades resulted from an increase in prey availability brought on by the decrease in baleen whale stocks. We question this tenet and present evidence to support the hypothesis that these increases are due to a gradual decrease in the frequency of cold years with extensive winter sea ice cover resulting from environmental warming. Supporting data were derived from one of the first, major multidisciplinary winter expedition to the Scotia and Weddell seas; recent satellite images of ocean ice cover; and the analysis of long-term surface temperature records and penguin demography. Our observations indicate there is a need to pay close attention to environmental data in the management of Southern Ocean resources given the complexity of relating biological changes to ecological perturbations.

Introduction

Populations of many krill-eating, Southern Ocean predators have exhibited significant changes during the last four decades. Notable among these, have been increases in the abundance of Chinstrap Penguins (*Pygoscelis antarctica*), which breed mainly on the Antarctic Peninsula and islands of the Scotia Sea (Watson 1975). At many colonies, numbers have increased 6–10% per annum (Laws 1985), and at some localities fivefold increases have occurred in the last 20 years (Rootes 1988). Chinstraps have also expanded their range southward along the western side of the Antarctic Peninsula (Parmelee and Parmelee 1987; Poncet and Poncet 1987) into areas historically dominated by the closely related adelic Penguin (*P. adeliae*; Fig. 1). A central tenet of Antarctic ecology ex-

plains these population changes in terms of a presumed increase in food availability that resulted from the decrease in baleen whale stocks due to commercial whaling (Sladen 1964; Emison 1968; Conroy 1975; Croxall and Kirkwood 1979; Croxall and Prince 1979; Croxall et al. 1984). This tenet is based on the fact that the dominant component in the summer diets of both Chinstraps and whales is the Antarctic krill (*Euphausia superba*). Although this tenet has been widely accepted, the possible mechanism by which a decrease in whales could have led to an increase in Chinstraps has not been questioned (cf. Horwood 1980). Indeed, the long-standing view has simply been that whaling led to a "krill surplus" that was used by krill-eating predators when competitive release altered the existing patterns of consumption (Laws 1985).

Although this whale reduction hypothesis has clearly been useful in guiding research on trophic interactions in the Southern Oceans, it is now apparent that increases in Chinstrap populations have not been mirrored by their sympatric, most closely related congener, the Adelic Penguin. Adelines share a significant portion of their range on the Antarctic peninsula and islands of the Scotia Arc with Chinstraps (Watson 1975). Alike in size and general appearance, both exhibit broad ecological similarities, not the least of which is a predominance of krill in their summer diets (Volkman et al. 1980; Trivelpiece et al. 1987, 1990; Trivelpiece and Trivelpiece 1990). Yet, when compared to Chinstraps, population increases in Adelines have not been as substantial, and at many sites appear to represent nothing more than recovery after human disturbance and exploitation (Poncet and Poncet 1987). Adelines, in fact, have declined noticeably at several localities on the Antarctic Peninsula, a change considered "unexplainable" by Poncet and Poncet (1987). This raises an interesting challenge to the whale reduction hypothesis: If the decrease of baleen whale stocks actually led to a krill surplus, why have populations of the ecologically similar Adelines residing in the same geographical areas shown such different responses?

Here we propose that the answer to this question does not rest with the idea of a krill surplus. Instead, we suggest

Key point: If the decimation of baleen whale populations did in fact lead to a "krill surplus", why were krill-dependent, top predator populations exhibiting such dichotomous trends?

"...the day bird people have something to tell us about climate warming is perhaps the day logic in climate science is abandoned..."

Anonymous Reviewer, Nature

"...a paper that creates this kind of controversy should be positive for science and the journal..."


G. Hempel, Editor, Pol. Biol.

* Current address: Polar Oceans Research Group, Department of Oceanography, Old Dominion University, Norfolk, VA 23529, USA.


** Present address: W.R. Fraser, ODU Central States Office 830 Hunt Farm Rd., Long Lake, MN 55356, USA

Offprint requests to: W.R. Fraser


The central hypothesis when the LTER began was that sea ice timing and magnitude structure the productivity and composition of the Antarctic ecosystem. The ice dynamics are driven by large-scale interactions of the atmosphere and ocean.




Winter 2007




Summer 2007







Temperature Trends (degrees C per year)



Feb 1979


Feb 1999



Sea Ice Concentration (%)

10 year analysis annual trends


Annual Rate of Sea Ice Concentration change (%)
1978–2008



ice decline


The WAP peninsula is experiencing the largest winter warming on Earth

Mean Winter Temperatures




Black is British Faraday & Ukraine Vernadsky Station
Red is US Palmer Station

Air temperature increases over the peninsula



Sea ice duration drops





The WAP peninsula is experiencing the largest winter warming on Earth




The WAP peninsula is experiencing the largest winter warming on Earth

Larson-B ice shelf after its collapse
Thanks to BAS & A. Clarke






In 2008 the Wilkens Ice Sheet followed the Larson Ice Shelf and began to collapse





Melt pools on surface of King George VI Sound


(from a BAS twin otter, January 2004)



Seasonal ice has declined over the few decades resulting to a climate migration to the South



Seasonal ice has declined over the few decades resulting to a climate migration to the South



Sea ice data courtesy of E. Chapman

**Warmer
moister**

*A climate gradient along the peninsula;
Warm, moist maritime conditions migrating south*

**Colder
drier**

Key Implications:
Regional shifts in the sea ice has major ecological implications

1990







Palmer Station in the present




photo by Bill Fraser


Plants at Palmer Station, the greening of Antarctica



Heat input from Antarctic Circumpolar Current (ACC - world's largest ocean current = ~30,000 Niagara Falls). The heat is driven onto the shelf by intensification of upwelling-favorable winds.





Heat input from Antarctic Circumpolar Current (ACC - world's largest ocean current = ~30,000 Niagara Falls). The heat is driven onto the shelf by intensification of upwelling-favorable winds.




Heat input from Antarctic Circumpolar Current (ACC - world's largest ocean current = ~30,000 Niagara Falls). The heat is driven onto the shelf by intensification of upwelling-favorable winds.

The WAP is the only location in the Antarctic where the ACC is adjacent to the shelf break. The ACC is Antarctica's warmest water







thanks to Doug Martinson



thanks to Doug Martinson

Upwelling favorable winds result in Ekman mass transport offshore






thanks to Doug Martinson

Heating on the WAP is driven by circulation and intrusion of the of the ACC onto the WAP continental shelf.


Meredith and King 2005



Summer sea surface temperature increasing +0.4 C/year

Evidence of solar ocean warming

Martinson et al. 2008





Ocean heat content increasing on the continental shelf of the WAP

Evidence of increased upwelling of Circumpolar Deep Waters


Thanks to
Doug
Martinson

Interannual variability is complex due to interacting influence of the Southern Annular Mode and El Nino/La Nina










Palmer time series: Phytoplankton show large interannual variability




What regulates phytoplankton blooms in this region?






Huang & Bender

The decadal changes have resulted changes in the phytoplankton




Montes Hugo et al. Science 2009




The changes driven by a decline in sea ice, wind and sun






15 year time
series of radio-
carbon
measurements
also suggest a
North & South
gradient

When chlorophyll is high, phytoplankton cells are big and are largely diatoms





Montes Hugo et al. 2009

Change in biomass impact on biogeochemistry





Who will dominate the warmer WAP?


● Prymnesiophytes □ Diatoms ■ Cryptophytes




Moline et al. GBC 2004



Moline et al.
GBC 2004




Moline et al.
GBC 2004



Corethron criophilum


Palmer Cryptophytes --> $8 \pm 2\mu\text{m}$

SEM Micrographs from McMinn and Hodgson 1993



100 μm


Thalassiosira antarctica



10 μm

Cryptomonas cryophila

A general feature in the warming WAP?



Location

South Shetland Islands

Weddell-Scotia-
Bellingshausen
Confluence Areas

Ellis Fjord

Bransfield Strait

Historical Data


Anvers Island
Signy Island

Reference

V illafañe et al., 1995;
Kang, S-H et al., 1997;
Kang, J-S et al., 1997
Lancelot et al., 1991;
Nothig et al., 1991
Tréguer et al., 1991;
Buma, 1992;
Mura et al., 1995;
Kang and Lee, 1995;
Aristegui et al., 1996
McMinn and Hodgson, 1993
Kang and Lee, 1995;
Kang et al., 1995


Krebs, 1983
Whitaker, 1982

CO₂ uptake varies with phytoplankton community




Montes Hugo in prep

Zooplankton are dominated by krill or salps




Krill greatest biomass of any animal on earth





From *Loeb et al.*, 1997






Is there an impact on higher trophic levels?






One focus idea of the LTER is testing, is that system is undergoing climate migration. We have structured sampling around the major Adelie penguin breeding areas along the peninsula.


Summer
foraging
areas for Adelie
penguins



Winter
foraging
areas for Adelie
penguins





To be expanded by NASA
grant awarded in Dec.



Anvers Island





Changing diets for the Adelie penguins



**Warmer
moister**




*A climate gradient along
the peninsula;
Warm, moist maritime
conditions migrating
south*

**Colder
drier**





Changing diets for the Adelie penguins

1994-present






Changing diets for the Adelie penguins


1994-present



1995-present








If that was not enough, warmer temps leads to more moisture and more snow. Breeding failure.....




BREEDING POPULATION CHANGE (%)




Results suggests the pelagic niche for the Adelie is changing

Satellite Defined Adelie Penguin Defined Environment (red) 2000–2003




Satellite Defined Adelie Penguin Defined Environment (red) 2005–2008



Adelie penguins
Weddell seals

Chinstrap penguins
Gentoo penguins
Fur seals
Elephant seals






Old Day Communication



HAM Operator Coms Palmer Station 1988



Brave New Day





RV Gould




Rutgers COOLroom




Enhanced productivity is associated with the warm upwelled water








Temperature from ru05's 01/24/09 - 01/29/09 deployment with overlay of 2009 penguin UD shown at 50% and 90% confidence intervals





Chlorophyll from ru05's 01/24/09 - 01/29/09 deployment with overlay of 2009 penguin UD shown at 50% and 90% confidence intervals





OOI planning & prosecution cyberinfrastructure



OOI planning & prosecution cyberinfrastructure



OOI planning & prosecution cyberinfrastructure



Conclusions:

Minor variations in the ocean state can have profound impacts on polar ecosystems

These profound changes are occurring in many polar oceans, changes appear to be accelerating

New technologies offer a mode to study and understand these changes, so it is time hopefully speed up our uphill trek to quantitative understanding, animals will help show us the way





RU-COOL Glider Fleet: Antarctic Deployments: 2007 – 2011

