



















# Oceans are complex and central to the Earth system













Despite heroic efforts, the ocean is chronically under-sampled. We need a permanent sustained presence making SPATIAL time series.





**Observation Lab** 







RUTGERS

Coastal Ocean Observation Lab

## **Rutgers University - Coastal Ocean Observation Lab**






**CODAR Network** 



L-Band & X-Band Satellite Receivers




Glider Fleet




3-D Nowcasts & Forecasts



#### **MARACOOS** - International Constellation of Satellites – Since 1992





## Mid-Atlantic Bight HF Radar Network – Since 1997






#### **MARACOOS - Autonomous Underwater Gliders – Since 1998**








**RU-COOL** Deployments by Year

Year	Number	Distance (km)	Days
2012	19	5054.31	290.3
2011	37	11975.53	624
2010	46	17003.36	876.6
2009	30	18114.34	761.1
2008	28	16642.28	604.7
2007	27	10277.99	462.9
2006	38	9745.88	538
2005	16	4113.87	207.1
Totals	241	92927.56	4364.8

Includes all Global deployments


- Since 2005: 241 Global deployments totaling 92,927 km over 4365 days
- Since 2005: 149 Mid-Atlantic deployments totaling 52,187 km over 2628 days








silbo Transect 01: 2012-07-24 12:46 - 2013-03-21 09:42 GMT

Models improved with in situ data via data assimilation











The Miami Herald > Weather >

# Hurricanes

Sunday, 02.19.12 Welcome Guest •



Miami-Dade | Broward | Keys | |

Posted on Friday, 09.02.11

HURRICANE SEASON

### Intensity remain



The National Hurricane Cer it wasn't the first time — bu the increasing standards of

BY CURTIS MORGAN
MORGAN@MIAMIHERALD.COM

#### The New Hork Times

U.S.

WORLD U.S. N.Y. / REGION BUSINESS TECHNOLOGY SCIENCE HEALTH SPORTS OPINION

POLITICS EDUCATION BAY AREA CHICAGO TEXAS



#### Challenges in Predicting the Intensity of Storms



Scientists say that it is much easier to accurately predict what path a hurricane will take.


By HENRY FOUNTAIN

Published: August 27, 2011




#### **Hurricane Irene**

#### Long Range Radar Network Sea Surface Currents 2011082717 GMT




#### 39.5N 73W Surface Current Time Series



#### **Hurricane Irene**


#### **Direct Wind Forcing**


# Mean Wavelet Power between Period 20 h and 30 h 2011-08-23 00:00:00




#### **Inertial Response**

Mean Wavelet Power between Period 16 h and 20 h 2011-08-23 00:00:00








WARM: RTG+SPoRT(8/27)


Init: 2011-08-27\_00:00:00

RTG,SPoRT(9/06)+AVHRR(8/31) Init: 2011-08-28\_06:00:00


#### 28/0600→29/0600

27/1200<del>→</del>28/0600






RUTGERS



#### WARM (RTG only)


#### COLD "Update" (AVHRR)







1. '.	NHC Forecast	only)	Warm (RTG only, OML Model)	Cold "Update" (AVHRR)
27/1200	5	-17.22	-17.23	-6.17
27/1800	10	4.1	4.2	5.88
28/0000	10	1.39	-2.14	3.96
28/0600	5	-1.2	-1.04	-1.21
28/1200	15	2.39	4.79	0.5
28/1800	15	4.97	3.51	-2.67
29/0000	15	3.62	1.93	-0.89
29/0600	10	10.48	9.84	4.52
<b>Sum of Squares</b>	800	457	452	118
RMSE	9.43	7.13	7.09	3.61







# Temperature






# Salinity







### Temperature profile




# Temperature change with depth (dT/dz)






Mixed layer depth = max(dT/dz)

Transition layer= dT/dz< 0.01












WBL grows and mixes from the bottom up


Winds mix from the top down














#### We are using the WAP as zone to study how rapid warming can shift the food web






Heat engine melting the WAP is the ACC


ACC upwelling is rapid flushing events (time scale of weeks)





track distance (km)


 Glider depth-averaged currents: agree well with moorings + model!




Thanks Lamont team




- Glider depth-averaged currents: agree well with moorings + model!
- Glider deep water (> 220 db) temperatures: episodic warming!










- •After about 3 days at the fence, we begin tracking.
- •Divert north/south to wait for feature to catch up if we exit  $(v_g \sim 3v_{bg})$ .
- •Follow feature for about 6 days.




## There are indications in changes in the primary production

## Remote Sensing (Decadal changes)



#### Montes-Hugo et al. 2009

### Ship Observed (Decadal changes)




Vernet et al. 2008






Antarctic krill




Antarctic Salp

## There are indications in changes in the zooplankton communities





Loeb et al. 1997







# **Adaptive Sampling of Penguin Habitats**
















































## Weather











## An ensemble of different numerical models run by different laboratories






# A fleet of four gliders October 30 – November 20, 2009




4 Deployments (3 Rutgers, I University of Delaware)

77 in-water days


1608 km flown



# COMBINING THE FIELD ASSETTS WITH OCEAN FORECASTS FOR PLANNING AND PROSECUTION EFFORTS




- Known constraints (slow 0.5 knot, Battery, shipping lanes)
- Uncertain constraints (time-varying 3D currents)
- Operate autonomously & re-plan daily




- From A to B in the shortest time
- Follow a time-varying feature (shelfslope salinity intrusion)





#### PLANNING FOR THE PRESENT





#### PLANNING IN THE FUTURE

